ANSWERS

Chapter 9

9.1 1.8

9.2 (a) From the given graph for a stress of 150×10^6 N m^{-2} the strain is 0.002

(b) Approximate yield strength of the material is 3×10^8 N m^{-2}

9.3 (a) Material A

(b) Strength of a material is determined by the amount of stress required to cause fracture: material A is stronger than material B.

9.4 (a) False (b) True

9.5 1.5×10^{-4} m (steel); 1.3×10^{-4} m (brass)

9.6 Deflection = 4×10^{-6} m

9.7 2.8×10^{-6}

9.8 0.127

9.9 7.07×10^4 N

9.10 $D_{copper}/D_{iron} = 1.25$

9.11 1.539×10^{-4} m

9.12 2.026×10^9 Pa

9.13 1.034×10^3 kg/m^3

9.14 0.0027

9.15 0.058 cm^3

9.16 2.2×10^6 N/m^2

9.17 Pressure at the tip of anvil is 2.5×10^{11} Pa

9.18 (a) 0.7 m (b) 0.43 m from steel wire

9.19 Approximately 0.01 m

9.20 260 kN

9.21 $2.51 \times 10^{-4}\ m^3$

Chapter 10

10.3 (a) decreases (b) η of gases increases, η of liquid decreases with temperature (c) shear strain, rate of shear strain (d) conservation of mass, Bernoulli's equation (e) greater.

10.5 6.2×10^6 Pa

10.6 10.5 m

10.7 Pressure at that depth in the sea is about 3×10^7 Pa. The structure is suitable since it can withstand far greater pressure or stress.

10.8 6.92×10^5 Pa

10.9 0.800

10.10 Mercury will rise in the arm containing spirit; the difference in levels of mercury will be 0.221 cm.

10.11 No, Bernoulli's principle applies to streamline flow only.

10.12 No, unless the atmospheric pressures at the two points where Bernoulli's equation is applied are significantly different.

10.13 9.8×10^2 Pa (The Reynolds number is about 0.3 so the flow is laminar).

10.14 1.5×10^3 N

10.15 Fig (a) is incorrect [Reason: at a constriction (i.e. where the area of cross-section of the tube is smaller), flow speed is larger due to mass conservation. Consequently pressure there is smaller according to Bernoulli's equation. We assume the fluid to be incompressible].

10.16 0.64 m s^{-1}

10.17 2.5×10^{-2} N m^{-1}

10.18 4.5×10^{-2} N for (b) and (c), the same as in (a).

10.19 Excess pressure = 310 Pa, total pressure = 1.0131×10^5 Pa. However, since data are correct to three significant figures, we should write total pressure inside the drop as 1.01×10^5 Pa.

© BSTBPC WEB COPY NOT TO BE PUBLISHED

10.20 Excess pressure inside the soap bubble = 20.0 Pa; excess pressure inside the air bubble in soap solution = 10.0 Pa. Outside pressure for air bubble = $1.01 \times 10^5 + 0.4 \times 10^3 \times 9.8 \times 1.2 = 1.06 \times 10^5$ Pa. The excess pressure is so small that up to three significant figures, total pressure inside the air bubble is 1.06×10^5 Pa.

10.21 55 N (Note, the base area does not affect the answer)

10.22 (a) absolute pressure = 96 cm of Hg; gauge pressure = 20 cm of Hg for (a), absolute pressure = 58 cm of Hg, gauge pressure = -18 cm of Hg for (b); (b) mercury would rise in the left limb such that the difference in its levels in the two limbs becomes19 cm.

10.23 Pressure (and therefore force) on the two equal base areas are identical. But force is exerted by water on the sides of the vessels also, which has a nonzero vertical component when the sides of the vessel are not perfectly normal to the base. This net vertical component of force by water on sides of the vessel is greater for the first vessel than the second. Hence the vessels weigh different even when the force on the base is the same in the two cases.

10.24 0.2 m

10.25 (a) The pressure drop is greater (b) More important with increasing flow velocity.

10.26 (a) 0.98 m s^{-1}; (b) 1.24×10^{-5} m^3 s^{-1}

10.27 4393 kg

10.28 5.8 cm s^{-1}, 3.9×10^{-10} N

10.29 5.34 mm

10.30 For the first bore, pressure difference (between the concave and convex side) = $2 \times 7.3 \times 10^{-2} / 3 \times 10^{-3} = 48.7$ Pa. Similarly for the second bore, pressure difference = 97.3 Pa. Consequently, the level difference in the two bores is $[48.7 / (10^3 \times 9.8)]$ m = 5.0 mm.

The level in the narrower bore is higher. (Note, for zero angle of contact, the radius of the meniscus equals radius of the bore. The concave side of the surface in each bore is at 1 atm).

10.31 (b) 8 km. If we consider the variation of g with altitude the height is somewhat more, about 8.2 km.

Chapter 11

11.1 Neon: – 248.58 °C = – 415.44 °F;

CO_2: – 56.60 °C = – 69.88 °F

(use $t_F = \frac{9}{5} t_c + 32$)

11.2 $T_A = (4/7)\ T_B$

11.3 384.8 K

11.4 (a) Triple-point has a *unique* temperature; fusion point and boiling point temperatures depend on pressure; (b) The other fixed point is the absolute zero itself; (c) Triple-point is 0.01°C, not 0 °C; (d) 491.69.

11.5 (a) T_A = 392.69 K, T_B = 391.98 K; (b) The discrepancy arises because the gases are not perfectly ideal. To reduce the discrepancy, readings should be taken for lower and lower pressures and the plot between temperature measured versus absolute pressure of the gas at triple point should be extrapolated to obtain temperature in the limit pressure tends to zero, when the gases approach ideal gas behaviour.

11.6 Actual length of the rod at 45.0 °C = (63.0 + 0.0136) cm = 63.0136 cm. (However, we should say that change in length up to three significant figures is 0.0136 cm, but the total length is 63.0 cm, up to three significant places. Length of the same rod at 27.0 °C = 63.0 cm.

11.7 When the shaft is cooled to temperature – 69°C the wheel can slip on the shaft.

11.8 The diameter increases by an amount = 1.44×10^{-2} cm.

11.9 3.8×10^2 N

11.10 Since the ends of the combined rod are not clamped, each rod expands freely.

Δl_{brass} = 0.21 cm, Δl_{steel} = 0.126 cm = 0.13 cm

Total change in length = 0.34 cm. No 'thermal stress' is developed at the junction since the rods freely expand.

11.11 0.0147 = 1.5×10^{-2}

11.12 103 °C

11.13 1.5 kg

11.14 0.43 J g^{-1} K^{-1} ; smaller

11.15 The gases are diatomic, and have other degrees of freedom (i.e. have other modes of motion) possible besides the translational degrees of freedom. To raise the temperature of the gas by a certain amount, heat is to be supplied to increase the average energy of all the modes. Consequently, molar specific heat of diatomic gases is more than that of monatomic gases. It can be shown that if only rotational modes of motion are considered, the molar specific heat of diatomic gases is nearly (5/2) R which agrees with the observations for all the gases listed in the table, except chlorine. The higher value of molar specific heat of chlorine indicates that besides rotational modes, vibrational modes are also present in chlorine at room temperature.

11.16 (a) At the triple point temperature = – 56.6 °C and pressure = 5.11 atm.

(b) Both the boiling point and freezing point of CO_2 decrease if pressure decreases.

(c) The critical temperature and pressure of CO_2 are 31.1 °C and 73.0 atm respectively. Above this temperature, CO_2 will not liquefy even if compressed to high pressures.

(d) (a) vapour (b) solid (c) liquid

11.17 (a) No, vapour condenses to solid directly.

(b) It condenses to solid directly without passing through the liquid phase.

(c) It turns to liquid phase and then to vapour phase. The fusion and boiling points are where the horizontal line on P–T diagram at the constant pressure of 10 atm intersects the fusion and vaporisation curves.

(d) It will not exhibit any clear transition to the liquid phase, but will depart more and more from ideal gas behaviour as its pressure increases.

© BSTBPC WEB COPY NOT TO BE PUBLISHED

11.18 4.3 g/min

11.19 3.7 kg

11.20 238 °C

11.22 9 min

Chapter 12

12.1 16 g per min

12.2 934 J

12.4 2.64

12.5 16.9 J

12.6 (a) 0.5 atm (b) zero (c) zero (assuming the gas to be ideal) (d) No, since the process (called free expansion) is rapid and cannot be controlled. The intermediate states are non-equilibrium states and do not satisfy the gas equation. In due course, the gas does return to an equilibrium state.

12.7 15%, 3.1×10^{9} J

12.8 25 W

12.9 450 J

12.10 10.4

Chapter 13

13.1 4×10^{-4}

13.3 (a) The dotted plot corresponds to 'ideal' gas behaviour; (b) $T_1 > T_2$; (c) 0.26 J K^{-1}; (d) No, 6.3×10^{-5} kg of H_2 would yield the same value

13.4 0.14 kg

13.5 5.3×10^{-6} m^3

13.6 6.10×10^{26}

13.7 (a) 6.2×10^{-21} J (b) 1.24×10^{-19} J (c) 2.1×10^{-16} J

13.8 Yes, according to Avogadro's law. No, v_{rms} is largest for the lightest of the three gases; neon.

13.9 2.52×10^{3} K

13.10 Use the formula for mean free path :

$$\bar{l} = \frac{1}{\sqrt{2}\pi n d^2}$$

where d is the diameter of a molecule. For the given pressure and temperature $N/V = 5.10 \times 10^{25}$ m^{-3} and $= 1.0 \times 10^{-7}$ m. $v_{rms} = 5.1 \times 10^2$ m s^{-1}.

collisional frequency = $\frac{v_{rms}}{\bar{l}} = 5.1 \times 10^9$ s^{-1}. Time taken for the collision = $d / v_{rms} = 4 \times 10^{-13}$ s.

Time taken between successive collisions = $1 / v_{rms} = 2 \times 10^{-10}$ s. Thus the time taken between successive collisions is 500 times the time taken for a collision. Thus a molecule in a gas moves essentially free for most of the time.

13.11 Nearly 24 cm of mercury flows out, and the remaining 52 cm of mercury thread plus the 48 cm of air above it remain in equilibrium with the outside atmospheric pressure (We assume there is no change in temperature throughout).

13.12 Oxygen

13.14 Carbon[1.29 Å]; Gold [1.59 Å]; Liquid Nitrogen [1.77 Å]; Lithium [1.73 Å]; Liquid fluorine[1.88 Å]

Chapter 14

14.1 (b), (c)

14.2 (b) and (c): SHM; (a) and (d) represent periodic but not SHM [A polyatomic molecule has a number of natural frequencies; so in general, its vibration is a superposition of SHM's of a number of different frequencies. This superposition is periodic but not SHM].

14.3 (b) and (d) are periodic, each with a period of 2 s; (a) and (c) are not periodic. [Note in (c), repetition of merely one position is not enough for motion to be periodic; the entire motion during one period must be repeated successively].

14.4 (a) Simple harmonic, $T = (2\pi/\omega)$; (b) periodic, $T = (2\pi/\omega)$ but not simple harmonic; (c) simple harmonic, $T = (\pi/\omega)$; (d) periodic, $T = (2\pi/\omega)$ but not simple harmonic; (e) non-periodic; (f) non-periodic (physically not acceptable as the function $\to \infty$ as $t \to \infty$.

14.5 (a) 0, +, + ; (b) 0, –, – ; (c) –, 0,0 ; (d) –, –, – ; (e) +, +, + ; (f) –, –, –.

14.6 (c) represents a simple harmonic motion.

14.7 A = $\sqrt{2}$ cm, $\phi = 7\pi/4$; B = $\sqrt{2}$ cm, $a = \pi/4$.

14.8 219 N

14.9 Frequency 3.2 s^{-1}; maximum acceleration of the mass 8.0 m s^{-2}; maximum speed of the mass 0.4 m s^{-1}.

14.10 (a) $x = 2 \sin 20t$
(b) $x = 2 \cos 20t$
(c) $x = -2 \cos 20t$

© BSTBPT WEB COPY, NOT TO BE PUBLISHED

where x is in cm. These functions differ neither in amplitude nor frequency. They differ in initial phase.

14.11 (a) $x = -3 \sin \pi t$ where x is in cm.

(b) $x = -2 \cos \frac{\pi}{2} t$ where x is in cm.

14.13 (a) F/k for both (a) and (b).

(b) $T = 2\pi \sqrt{\frac{m}{k}}$ for (a) and $2\pi \sqrt{\frac{m}{2k}}$ for (b)

14.14 100 m/min

14.15 8.4 s

14.16 (a) For a simple pendulum, k itself is proportional to m, so m cancels out.

(b) $\sin \theta < \theta$; if the restoring force, $mg \sin \theta$ is replaced by $mg\theta$, this amounts to effective reduction in g for large angles and hence an increase in time period T over that given by the formula $T = 2\pi \sqrt{\frac{l}{g}}$ where one assumes $\sin\theta = \theta$.

(c) Yes, the motion in the wristwatch depends on spring action and has nothing to do with acceleration due to gravity.

(d) Gravity disappears for a man under free fall, so frequency is zero.

14.17 $T = 2\pi \sqrt{\frac{l}{\sqrt{g^2 + v^4 / R^2}}}$. Hint: Effective acceleration due to gravity will get reduced due to radial acceleration v^2/R acting in the horizontal plane.

14.18 In equilibrium, weight of the cork equals the up thrust. When the cork is depressed by an amount x, the net upward force is $Ax\rho_l g$. Thus the force constant $k = A\rho_l g$.

Using $m = Ah\rho$, and $T = 2\pi \sqrt{\frac{m}{k}}$ one gets the given expression.

14.19 When both the ends are open to the atmosphere, and the difference in levels of the liquid in the two arms is h, the net force on the liquid column is $Ah\rho g$ where A is the area of cross-section of the tube and ρ is the density of the liquid. Since restoring force is proportional to h, motion is simple harmonic.

14.20 $T = 2\pi\sqrt{\frac{Vm}{Ba^2}}$ where B is the bulk modulus of air. For isothermal changes $B = P$.

14.21 (a) 5×10^4 N m^{-1}; (b) 1344.6 kg s^{-1}

14.22 Hint: Average K.E. = $\frac{1}{T}\int_0^T \frac{1}{2}mv^2 dt$; Average P.E.= $\frac{1}{T}\int_0^T \frac{1}{2}kx^2 dt$

14.23 Hint: The time period of a torsional pendulum is given by $T = 2\pi\sqrt{\frac{I}{\alpha}}$, where I is the moment of inertia about the axis of rotation. In our case $I = \frac{1}{2}MR^2$, where M is the mass of the disk and R its radius. Substituting the given values, $\alpha = 2.0$ N m rad^{-1}.

14.24 (a) $-5\pi^2$ m s^{-2}; 0; (b) $-3\pi^2$ m s^{-2}; 0.4π m s^{-1}; (c) 0 ; $0.5\,\pi$ m s^{-1}

14.25 $\sqrt{\left(x_0^2 + \frac{v_0^2}{\omega^2}\right)}$

Chapter 15

15.1 0.5 s

15.2 8.7 s

15.3 2.06×10^4 N

15.4 Assume ideal gas law: $P = \frac{\rho RT}{M}$, where ρ is the density, M is the molecular mass, and T is the temperature of the gas. This gives $v = \sqrt{\frac{\gamma RT}{M}}$. This shows that v is:

(a) Independent of pressure.

(b) Increases as $\sqrt{T}$.

(c) The molecular mass of water (18) is less than that of N_2 (28) and O_2 (32).

Therefore as humidity increases, the effective molecular mass of air decreases and hence v increases.

© BSTBPC WEB COPY NOT TO BE PUBLISHED

15.5 The converse is not true. An obvious requirement for an acceptable function for a travelling wave is that it should be finite everywhere and at all times. Only function (c) satisfies this condition, the remaining functions cannot possibly represent a travelling wave.

15.6 (a) 3.4×10^{-4} m (b) 1.49×10^{-3} m

15.7 4.1×10^{-4} m

15.8 (a) A travelling wave. It travels from right to left with a speed of 20 ms^{-1}.

(b) 3.0 cm, 5.7 Hz

(c) $\pi/4$

(d) 3.5 m

15.9 All the graphs are sinusoidal. They have same amplitude and frequency, but different initial phases.

15.10 (a) 6.4 π rad

(b) 0.8 π rad

(c) π rad

(d) $(\pi/2)$ rad

15.11 (a) Stationary wave

(b) l = 3 m, n = 60 Hz, and v = 180 m s^{-1} for each wave

(c) 648 N

15.12 (a) All the points except the nodes on the string have the same frequency and phase, but not the same amplitude.

(b) 0.042 m

15.13 (a) Stationary wave.

(b) Unacceptable function for any wave.

(c) Travelling harmonic wave.

(d) Superposition of two stationary waves.

15.14 (a) 79 m s^{-1}

(b) 248 N

15.15 347 m s^{-1}

Hint : $v_n = \frac{(2n-1)v}{4l}$; n = 1,2,3,....for a pipe with one end closed

15.16 5.06 km s^{-1}

15.17 First harmonic (fundamental); No.

15.18 318 Hz

15.20 (i) (a) 412 Hz, (b) 389 Hz, (ii) 340 m s^{-1} in each case.

15.21 400 Hz, 0.875 m, 350 m s^{-1}. No, because in this case, with respect to the medium, both the observer and the source are in motion.

15.22 (a) 1.666 cm, 87.75 cm s^{-1}; No, the velocity of wave propagation is – 24 m s^{-1}

(b) All points at distances of $n\lambda$ ($n = \pm1, \pm2, \pm3,....$) where λ = 12.6 m from the point x = 1 cm.

15.23 (a) The pulse does not have a definite wavelength or frequency, but has a definite speed of propagation (in a non-dispersive medium).

(b) No

15.24 $y = 0.05 \sin(\omega t - kx)$; here $\omega = 1.61 \times 10^3$ s^{-1}, $k = 4.84$ m^{-1}; x and y are in m.

15.25 45.9 kHz

15.26 1920 km

15.27 42.47 kHz

© BSTBPC WEB COPY ,NOT TO BE PUBLISHED

BIBLIOGRAPHY

TEXTBOOKS

For additional reading on the topics covered in this book, you may like to consult one or more of the following books. Some of these books however are more advanced and contain many more topics than this book.

1. **Ordinary Level Physics**, A.F. Abbott, Arnold-Heinemann (1984).
2. **Advanced Level Physics**, M. Nelkon and P. Parker, 6th Edition Arnold-Heinemann (1987).
3. **Advanced Physics**, Tom Duncan, John Murray (2000).
4. **Fundamentals of Physics**, David Halliday, Robert Resnick and Jearl Walker, 7th Edition John Wily (2004).
5. **University Physics**, H.D. Young, M.W. Zemansky and F.W. Sears, Narosa Pub. House (1982).
6. **Problems in Elementary Physics**, B. Bukhovtsa, V. Krivchenkov, G. Myakishev and V. Shalnov, MIR Publishers, (1971).
7. **Lectures on Physics** (3 volumes), R.P. Feynman, Addision – Wesley (1965).
8. **Berkeley Physics Course** (5 volumes) McGraw Hill (1965).
   a. Vol. 1 – Mechanics: (Kittel, Knight and Ruderman)
   b. Vol. 2 – Electricity and Magnetism (E.M. Purcell)
   c. Vol. 3 – Waves and Oscillations (Frank S. Craw-ford)
   d. Vol. 4 – Quantum Physics (Wichmann)
   e. Vol. 5 – Statistical Physics (F. Reif)
9. **Fundamental University Physics**, M. Alonso and E. J. Finn, Addison – Wesley (1967).
10. **College Physics**, R.L. Weber, K.V. Manning, M.W. White and G.A. Weygand, Tata McGraw Hill (1977).
11. **Physics**: **Foundations and Frontiers**, G. Gamow and J.M. Cleveland, Tata McGraw Hill (1978).
12. **Physics for the Inquiring Mind**, E.M. Rogers, Princeton University Press (1960)
13. **PSSC Physics Course**, DC Heath and Co. (1965) Indian Edition, NCERT (1967)
14. **Physics Advanced Level**, Jim Breithampt, Stanley Thornes Publishers (2000).
15. **Physics,** Patrick Fullick, Heinemann (2000).

16. **Conceptual Physics,** Paul G. Hewitt, Addision-Wesley (1998).
17. **College Physics,** Raymond A. Serway and Jerry S. Faughn, Harcourt Brace and Co. (1999).
18. **University Physics,** Harris Benson, John Wiley (1996).
19. **University Physics,** William P. Crummet and Arthur B. Western, Wm.C. Brown (1994).
20. **General Physics,** Morton M. Sternheim and Joseph W. Kane, John Wiley (1988).
21. **Physics,** Hans C. Ohanian, W.W. Norton (1989).
22. **Advanced Physics,** Keith Gibbs, Cambridge University Press(1996).
23. **Understanding Basic Mechanics,** F. Reif, John Wiley (1995).
24. **College Physics,** Jerry D. Wilson and Anthony J. Buffa, Prentice-Hall (1997).
25. **Senior Physics, Part – I,** I.K. Kikoin and A.K. Kikoin, Mir Publishers (1987).
26. **Senior Physics, Part – II,** B. Bekhovtsev, Mir Publishers (1988).
27. **Understanding Physics,** K. Cummings, Patrick J. Cooney, Priscilla W. Laws and Edward F. Redish, John Wiley (2005)
28. **Essentials of Physics,** John D. Cutnell and Kenneth W. Johnson, John Wiley (2005)

GENERAL BOOKS

For instructive and entertaining general reading on science, you may like to read some of the following books. Remember however, that many of these books are written at a level far beyond the level of the present book.

1. **Mr. Tompkins** in paperback, G. Gamow, Cambridge University Press (1967).
2. **The Universe and Dr. Einstein**, C. Barnett, Time Inc. New York (1962).
3. **Thirty years that Shook Physics**, G. Gamow, Double Day, New York (1966).
4. **Surely You're Joking, Mr. Feynman**, R.P. Feynman, Bantam books (1986).
5. **One, Two, Three... Infinity**, G. Gamow, Viking Inc. (1961).
6. **The Meaning of Relativity**, A. Einstein, (Indian Edition) Oxford and IBH Pub. Co (1965).
7. **Atomic Theory and the Description of Nature**, Niels Bohr, Cambridge (1934).
8. **The Physical Principles of Quantum Theory**, W. Heisenberg, University of Chicago Press (1930).
9. **The Physics- Astronomy Frontier**, F. Hoyle and J.V. Narlikar, W.H. Freeman (1980).
10. **The Flying Circus of Physics with Answer**, J. Walker, John Wiley and Sons (1977).
11. **Physics for Everyone** (series), L.D. Landau and A.I. Kitaigorodski, MIR Publisher (1978).
    Book 1: Physical Bodies
    Book 2: Molecules
    Book 3: Electrons
    Book 4: Photons and Nuclei.
12. **Physics can be Fun**, Y. Perelman, MIR Publishers (1986).
13. **Power of Ten**, Philip Morrison and Eames, W.H. Freeman (1985).
14. **Physics in your Kitchen Lab**., I.K. Kikoin, MIR Publishers (1985).
15. **How Things Work : The Physics of Everyday Life**, Louis A. Bloomfield, John Wiley (2005)
16. **Physics Matters : An Introduction to Conceptual Physics,** James Trefil and Robert M. Hazen, John Wiley (2004).

INDEX

A

Absolute scale temperature 276
Absolute zero 276
Acceleration (linear) 45
Acceleration due to gravity 49,189
Accuracy 22
Action-reaction 97
Addition of vectors 67
Adiabatic process 306, 307
Aerofoil 258
Air resistance 79
Amplitude 340, 368
Angle of contact 263, 264
Angstrom 21
Angular Acceleration 154
Angular displacement 338
Angular frequency 340, 369
Angular momentum 155
Angular velocity 152
Angular wave number 368
Antinodes 376
Archimedes Principle 251
Area expansion 277
Atmospheric pressure 249
Average acceleration 45, 74
Average speed 42
Average velocity 42
Avogardo's law 320

B

Banked road 104
Barometer 250
Beat frequency 338, 380, 381
Beats 379, 380
Bending of beam 240
Bernoulli's Principle 254
Blood pressure 272, 273
Boiling point 283
Boyle's law 321
Buckling 240
Bulk modulus 238
Buoyant force 251

C

Calorimeter 281
Capillary rise 264
Capillary waves 366
Carnot engine 311
Central forces 186
Centre of Gravity 161
Centre of mass 144
Centripetal acceleration 81
Centripetal force 104
Change of state 282
Charle's law 321
Chemical Energy 126
Circular motion 104
Clausius statement 310
Coefficient of area expansion 279
Coefficient of linear expansion 277
Coefficient of performance 309
Coefficient of static friction 101
Coefficient of viscosity 258
Coefficient of volume expansion 277
Cold reservoir 308
Collision 129
Collision in two dimensions 131
Compressibility 238
Compressions 364, 365, 371
Compressive stress 232, 239
Conduction 286
Conservation laws 12
Conservation of angular momentum 157, 173
Conservation of Mechanical Energy 121
Conservation of momentum 98
Conservative force 121
Constant acceleration 46,75
Contact force 100
Convection 289
Couple 159
Crest 367
Cyclic process 307

D

Dalton's law of partial pressure 321
Damped oscillations 351
Damped simple Harmonic motion 351
Damping constant 351
Damping force 351
Derived units 16
Detergent action 265
Diastolic pressure 273
Differential calculus 61
Dimensional analysis 32
Dimensions 31
Displacement vector 66
Displacement 40
Doppler effect 381, 382
Doppler shift 383
Driving frequency 354
Dynamics of rotational motion 169

E

Efficiency of heat engine 308
Elastic Collision 129
Elastic deformation 232, 234
Elastic limit 234
Elastic moduli 235
Elasticity 231
Elastomers 235
Electromagnetic force 8
Energy 117
Equality of vectors 66
Equation of continuity 253
Equilibrium of a particle 99
Equilibrium of Rigid body 158
Equilibrium position 337, 338, 349
Errors in measurement 22
Escape speed 193

F

First law of Thermodynamics 302
Fluid pressure 247
Force 94
Forced frequency 353
Forced oscillations 353
Fracture point 234
Free Fall 49
Free-body diagram 100
Frequency of periodic motion 338, 369
Friction 101
Fundamental Forces 6
Fundamental mode 377
Fusion 282

G

Gauge pressure 249
Geocentric model 183
Geostationary satellite 196
Gravitational constant 189
Gravitational Force 8, 192
Gravitational potential energy 191
Gravity waves 366

H

Harmonic frequency 377, 378
Harmonics 377, 378
Heat capacity 280
Heat engines 308
Heat pumps 308
Heat 274
Heliocentric model 183
Hertz 338
Hooke's law 234
Horizontal range 78
Hot reservoir 308
Hydraulic brakes 251, 252
Hydraulic lift 251, 252
Hydraulic machines 251
Hydraulic pressure 234
Hydraulic stress 234, 239
Hydrostatic paradox 249

I

Ideal gas equation 276
Ideal gas 321
Impulse 96
Inelastic collision 129
Initial phase angle 368
Instantaneous acceleration 74
Instantaneous speed 45
Instantaneous velocity 43
Interference 374
Internal energy 301, 325
Irreversible engine 311, 313
Irreversible processes 310
Isobaric process 306, 307
Isochoric process 306, 307
Isotherm 305
Isothermal process 306

K

Kelvin-Planck statement 310
Kepler's laws of planetary motion 184
Kinematics of Rotational Motion 167
Kinematics 39
Kinetic energy of rolling motion 174
Kinetic Energy 117
Kinetic interpretation of temperature 324
Kinetic theory of gases 323

L

Laminar flow 254, 260
Laplace correction 373

Latent heat of fusion 285
Latent heat of vaporisation 285
Latent heat 284
Law of cosine 72
Law of equipartition of energy 327
Law of Inertia 90
Law of sine 72
Linear expansion 277
Linear harmonic oscillator 345, 347
Linear momentum 155
Longitudinal strain 232
Longitudinal strain 232, 235
Longitudinal stress 232
Longitudinal Wave 366, 372

M

Magnus effect 257
Manometer 250
Mass Energy Equivalence 126
Maximum height of projectile 78
Maxwell Distribution 326
Mean free path 319, 330
Measurement of length 18
Measurement of mass 21
Measurement of temperature 275
Measurement of time 22
Melting point 282
Modes 376
Modulus of elasticity 234
Modulus of rigidity 238
Molar specific heat capacity at constant pressure 280, 304
Molar specific heat capacity at constant volume 280, 304
Molar specific heat capacity 280
Molecular nature of matter 318
Moment of Inertia 163
Momentum 93
Motion in a plane 72
Multiplication of vectors 67
Musical instruments 380

N

Natural frequency 354
Newton's first law of motion 91
Newton's Law of cooling 290
Newton's law of gravitation 185
Newton's second law of motion 93
Newton's third law of motion 96
Newtons' formula for speed of sound 372
Nodes 376
Normal Modes 376, 377, 379
Note 380, 381
Nuclear Energy 126
Null vector 68

O

Odd harmonics 378
Orbital velocity/speed 194
Order of magnitude 28
Oscillations 337
Oscillatory motion 337

P

Parallax method 18
Parallelogram law of addition of vectors 66
Pascal's law 248
Path length 40
Path of projectile 78
Periodic force 353
Periodic motion 337
Periodic time 337
Permanent set 234
Phase angle 340
Phase constant 340
Phase diagram 283
Pipe open at both ends 380
Pipe open at one end 379
Pitch 382
Plastic deformation 234
Plasticity 231
Polar satellite 196
Position vector and displacement 73
Potential energy of a spring 123
Potential energy 120
Power 128
Precession 143
Pressure gauge 249
Pressure of an ideal gas 323
Pressure pulse 366
Pressure 246
Principle of Conservation of Energy 128
Principle of moments 160
Progressive wave 366
Projectile motion 77
Projectile 77
Propagation constant 368
Pulse 365

Q

Quasi-static process 305, 306

R

Radial acceleration 344
Radiation 290
Radius of Gyration 164
Raman effect 11
Rarefactions 364, 365, 371
Ratio of specific heat capacities 329
Reaction time 51
Real gases 321
Rectilinear motion 39

Reductionism 2
Reflected wave 375
Reflection of waves 374
Refracted wave 375
Refrigerator 308
Regelation 282
Relative velocity in two dimensions 76
Relative velocity 51
Resolution of vectors 69
Resonance 354
Restoring force 232, 345, 364
Reversible engine 311, 312
Reversible processes 310
Reynolds number 260
Rigid body 141
Rolling motion 173
Root mean square speed 325
Rotation 142

S

S.H.M. (Simple Harmonic Motion) 339
Scalar-product 114
Scalars 65
Scientific Method 1
Second law of Thermodynamics 309
Shear modulus 238
Shearing strain 233
Shearing stress 232, 239
SI units 16
Significant figures 27
Simple pendulum 338, 349
Soap bubbles 264
Sonography 383
Sound 371
Specific heat capacity of Solids 303, 330
Specific heat capacity of Gases 328, 329
Specific heat capacity of Water 330
Specific heat capacity 280, 303
Speed of efflux 255
Speed of Sound 371, 372
Speed of Transverse wave on a stretched string 370, 371
Sphygmomanometer 273
Spring constant 348, 351
Standing waves 376
Stationary waves 378
Steady flow 253
Stethoscope 273
Stokes' law 259
Stopping distance 50
Strain 232
Streamline flow 253, 254
Streamline 253, 254
Stress 232
Stress-strain curve 234
Stretched string 370
Sublimation 284
Subtraction of vectors 67
Superposition principle 374
Surface energy 261
Surface tension 261
Symmetry 146
System of units 16
Systolic pressure 273

T

Temperature 275
Tensile strength 234
Tensile stress 232
Terminal velocity 260
Theorem of parallel axes 167
Theorem of perpendicular axes 165
Thermal conductivity 287
Thermal equilibrium 299
Thermal expansion 276
Thermal stress 279
Thermodynamic processes 305
Thermodynamic state variables 304
Thermodynamics 3, 298
Time of flight 78
Torque 154
Torricelli's Law 255, 256
Trade wind 289
Transmitted wave 375
Travelling wave 376
Triangle law of addition of vectors 66
Triple point 283
Trough 367
Tune 380
Turbulent flow 253, 254

U

Ultimate strength 234
Ultrasonic waves 383
Unification of Forces 10
Unified Atomic Mass Unit 21
Uniform circular motion 79
Uniform Motion 41
Uniformly accelerated motion 47
Unit vectors 70

V

Vane 351
Vaporisation 283
Vector-product 151
Vectors 66
Velocity amplitude 344
Venturi meter 256
Vibration 337
Viscosity 258
Volume expansion 277
Volume Strain 234

W

Wave equation 370
Wave length 368
Wave speed 370
Waves 363
Waxing and waning of sound 381
Weak nuclear force 9
Weightlessness 197
Work done by variable force 118
Work 116
Work-Energy Theorem 116
Working substance 308

Y

Yield Point 234
Yield strength 234
Young's modulus 235

Z

Zeroth law of Thermodynamics 300